

# Transmurale richtlijn fecale incontinentie en obstipatie

Claire Vrielink, Marieke Meeuwis, Quirine van der Weide-Warnaar en Mariska Slotman

Allen geregistreerd bekkenfysiotherapeut

## Literatuur

1. Meeuwis M, Slotman M, Vrielink C, van der Weide-Warnaar Q. Transmurale richtlijn fecale incontinentie en obstipatie. Amersfoort: NVFB; 2021.
2. Tanaz RF, Demtchouk V. Fecal incontinence [online]. MedScape. Beschikbaar via <https://emedicine.medscape.com/article/268674-overview>. Geraadpleegd 15 november 2020.
3. Koninklijk Nederlands Genootschap voor Fysiotherapie. KNGF-Evidence Statement Anale incontinentie. Amersfoort: KNGF; 2013.
4. Norton C, Cody JD. Biofeedback and/or sphincter exercises for the treatment of faecal incontinence in adults. Cochrane Database Syst Rev. 2012 Jul 11;7:CD002111.
5. ICI 6th edition 2017,
6. Skardoon GR, Khera AJ, Burgell ERE. Review article. Dyssynergic defaecation and biofeedback therapy in the pathophysiology and management of functional constipation. Alimentary Pharmacol Ther. 2017;46(4):410-23.
7. Belsey J, Greenfield S, Candy D, Geraint M. Systematic review, impact of constipation on quality of life in adults and children. Aliment Pharmacol Ther. 2010;31(9):938-49.
8. Rao SSC, Mudipalli RS, Stessman M, Zimmerman B. Investigation of the utility of colorectal function tests and Rome II criteria in dyssynergic defecation (Anismus). Neurogastroenterol Motil. 2004;16:589-96.
9. Diemel JM, van den Hurk APJM, Murris JWM, Pijpers MAM, Verheij AAA, Kurver MJ. NHG-Standaard obstipatie. Utrecht: Nederlandse Huisartsen Genootschap; 2010.
10. Rao SSC. A balancing view: Fecal incontinence: Test or treat empirically - Which strategy is best? Am J Gastroenterol. 2006;101(12), 2683-84.
11. Rao SSC, Rattanakot K, Patcharatrakul T. Diagnosis and management of chronic constipation in adults. Nat Rev Gastroenterol Hepatol. 2016;13(5):295-305.
12. Braekken IH, Majida M, Ellström Engh M, Bø K. Morphological changes after pelvic floor muscle training measured by 3-dimensional ultrasonography: a randomized controlled trial. Obstet Gynecol. 2010; 115(2 Pt 1):317-24.
13. Chiarione G, Whitehead WE, Pezza V, Morelli A, Bassotti G. Biofeedback is superior to laxatives for normal transit constipation due to pelvic floor dyssynergia. Gastroenterol. 2006;130(3):657-64.